

American earthworms (Oligochaeta) from north of the Rio Grande - a species checklist

by R.J. Blakemore YNU, Yokohama, Japan

December, 2006

Abstract This current review totals approximately 182 earthworm taxa in 12 families reported from America north of Mexico, i.e., USA & Canada, of which approximately 60 (ca. 33%) are exotic/introduced. All records are presented in an annotated checklist, but not all occurrences are verified and there may be some residual errors and/or omissions in this draft that, nevertheless, seems to be the most current and complete version available, at least online. Compare to Reynolds & Wetzel (2004) who claim 161 terrestrial species in 10 megadrile families from the same region, including just 45 exotics and many of these with superceded or obsolete names or family affiliations.

Introduction

The first (dubious) species described from America was *Hypogaeum hirtum* Savigny, 1820. Since then, the whole of North America (USA + Canada) – from the Arctic Ocean south to the Rio Grande but excluding offshore islands, had 161 valid species from 10 families of terrestrial oligochaetes reported by Reynolds & Wetzel (2004 – cited by Wetzel, 2005 here: inhs.uiuc.edu/~mjwetzels/AOGSMNP.PkChklist.html). Of these, 116 were considered native, and 45 introductions from other countries for which several of the names quoted had been superceded at least 20 years previously (eg. *Aporrectodea turgida*, "*Eisenia foetida* Savigny, 1826", *Amyntas hawayanus*, *A. diffringens* – see: inhs.uiuc.edu/~mjwetzels/TerrWoi.mjw.list.html May, 2005). Previously, Wetzel (2003) recognized 159 nominal species in 40 genera from 12 families and an earlier paper, by Reynolds (1995: 1) cited a study by "Reynolds & Wetzel, 1994" that check-listed 147 species from North America with about 30% exotic. In the same volume, now 10 years old, Fender (1995: 59) estimated the number of native North American taxa could be doubled, but it would take 5 years full time-work to describe the 100+ species and the results would be too big for conventional publication (cf. Gates, 1942). Schwert (1900) and Reynolds (1995: 10) listed 36-38 lumbricids; James (1995: 34) some 69 others, and Fender (1995: 53) mentioned 28 species (listed by Fender & McKey-Fender, 1900: 377), with perhaps another 80 awaiting description. These figures to 1995 give totals of about 133-135 described species and a potential count of 215 taxa. The current list is for species from continental North America (Canada and USA) north of the Rio Grande (i.e., north of Mexico) where the earthworm fauna is thought impoverished by extensive Wisconsin(i)an glaciations. However, in what is now northeastern USA, native Indians traditionally named Spring "Full Worm Moon" as its onset was heralded by earthworm activity (eg. surface casting, song birds feeding) preparing the earth for new growth.

Methods and Materials

This species list is updated from sources cited below and, with permission, from the **Polistes Foundation** website: usmo4.discoverlife.org/mp/20q?act=x_checklist&guide=Earthworms that lists 118 taxa (24 exotic : 94 native). Species from US territories, protectorates and zones of influence, e.g. Puerto Rico, Western Samoa, Hawaii, and Philippines are not included here (see separate checklists in this series).

Regarding the *Aporrectodea caliginosa* species-group complex *sensu* Blakemore (2002; 2006) [non *Allolobophora caliginosa* species-group, Gerard, 1964: 27; Sims & Gerard, 1985; 1999: 41, 56; nec *Allolobophora caliginosa* species-complex, Gates, 1972: 84, nec *Allolobophora trapezoides* species-group, Gates, 1972b: 1 nec *Aporrectodea trapezoides* species-complex, Easton, 1981: 41]; according to Sims & Gerard (1985; 1999: 44) some North American authors continue to incorrectly cite *Aporrectodea turgida* (Eisen, 1873) and/or *A. tuberculata* (Eisen, 1874) in preference to the prior *A. caliginosa* (Savigny, 1826) on the grounds that Savigny did not clearly define his species whereas Eisen's material permits adequate characterization (e.g. Gates, 1972; Reynolds, 1977; James, 1990: 381). Yet other authors, such as Schwert (1990: 355), acknowledge *trapezoides* and note that *turgida* is a junior synonym of *caliginosa* but also retain *tuberculata*. This latter strategy was supported by Blakemore (2002; 2006) and is also followed herein.

Results

Table 1. Estimated totals of native earthworm families, genera, and species in North America, north of Mexico (revised from Hendrix & Bohlen, 2002: tab. 2; Fender & McKey-Fender, 1990).

Family	Genus	Approx. No. Species
Ailoscolecidae	<i>Komarekiona</i> Gates, 1974	1
Lumbricidae	<i>Bimastos</i> Moore, 1893	10
	<i>Eisenoides</i> Gates, 1969	2
Lutodrilidae	<i>Lutodrilus</i> McMahan, 1976	1
Sparganophilidae	<i>Sparganophilus</i> Benham, 1892	12-13
Ocnerodrilidae	? <i>Ocnerodrilus</i>	1+
Acanthodrilidae	<i>Diplocardia</i> Garman, 1888	52-57
	? <i>Microscolex</i> Rosa, 1887	3?
Megascolecidae (all in tribe	<i>Arctiostrotus</i> McKey-Fender & Fender, 1982	7 (+2 undescribed?)
	<i>Argilophilus</i> Eisen, 1893	11 (+30 undescribed)
Argilophilini	<i>Chetcodrilus</i> Fender & McKey-Fender, 1990: 374	3 (+2 undescribed)
Fender &	<i>Drilocheira</i> Fender & McKey-Fender, 1990: 376	1 (+3 undescribed)
McKey-Fender, 1990)	<i>Driloleirus</i> Fender & McKey-Fender, 1990: 372	6 (+3 undescribed)
	<i>Kincaidodrilus</i> McKey-Fender, 1982	1
	<i>Macnabodrilus</i> Fender & McKey-Fender, 1990: 373	2
	<i>Nephralaxis</i> Fender & McKey-Fender, 1990: 375	2 (+7 undescribed)
	<i>Toutellus</i> Fender & McKey-Fender, 1990: 374	4 (+4 undescribed)
TOTALS	15-17	123-170

Species checklist

FAMILIES after Blakemore (2000); remarks and synonyms (syn.) in brackets, marked with "?" where there is some taxonomic uncertainty. Codes: * = exotic/introduced, - = native/endemic, # = uncertain affinities. For common exotics and cosmopolitan species, complete synonymies are found elsewhere (e.g. Sims & Gerard, 1985, 1999; Blakemore, 2000, 2002).

Family CRIODRILIDAE Vejdovsky, 1884 (Palearctic)

- *1. *Criodrilus lacuum* Hoffmeister, 1845 [syn. *Hydrilus ghaniae* Qiu & Bouché, 1998: 18, 198 synonymy of this monotypic genus was by Omodeo, Rota & Baha (2003: 6). Notes (from Blakemore, 2002): natural range from Iberian Peninsular to Pacific-coast of Siberia, and introduced into North America (plant pots, Baltimore)].

Family SPARGANOPHILIDAE Michaelsen, 1918 (Nearctic)

- 2. *Sparganophilus gatesi* Reynolds, 1980.
- 3. *Sparganophilus helenae* Reynolds, 1980.
- 4. *Sparganophilus komareki* Reynolds, 1980.
- 5. *Sparganophilus kristinae* Reynolds, 1980.
- 6. *Sparganophilus meansi* Reynolds, 1980.
- 7. *Sparganophilus pearsei libertiensis* Reynolds, 1980 [omitted by James (1995: 34)].
- 8. *Sparganophilus pearsei pearsei* Reynolds, 1975 [misspelt "pearsi" by James (1995: 34)].
- 9. *Sparganophilus pearsei sarasotae* Reynolds, 1980 [omitted by James (1995: 34)].
- 10. *Sparganophilus smithi* Eisen, 1896 (sometimes "smithii").
- 11. *Sparganophilus sonomae* Eisen, 1896.
- 12. *Sparganophilus tamesis* Benham, 1892 [syns. *S. eiseni* Smith, 1895 - misdated "Smith, 1980" by James (1995: 34) who attempts to maintain *S. eiseni* while at the same time overlooking *tamesis*, despite Sims & Gerard's (1985; 1999: 122) explanation of its synonymy; *Sparganophilus benhami* Eisen, 1896; *Sparganophilus benhami* var. *guatemalensis* Eisen, 1896; *Sparganophilus benhami* var. *carnea* Eisen, 1896; *Helodrilus elongatus* Friend, 1911; ?*Pelodrilus cuenoti* Tetry, 1934 [synonymy of *cuenoti* by Cernovitov reported in Gates (1972: 314), others from Sims & Gerard (1985, 1999: 120)]. Misspelt "tamesi" in Qiu & Bouché (1998: 178). Distribution is eastern North America through to Guatemala, introduced into Britain and France (Lorraine) - from Sims & Gerard (1999: 122)].
- 13. *Sparganophilus tennesseensis* Reynolds, 1977 [sometimes misspelt "tennessensis"].
- 14. *Sparganophilus wilmae* Reynolds, 1980.

Family AILOSCOLECIDAE (Pyrenees and southeast USA).

Note: Sims, (1980: 108) put American Komarekionidae Gates, 1974 in synonymy, cf. Reynolds & Cook, (1993).

- 15. *Komarekiona eatoni* Gates, 1974

Family HORMOGASTRIDAE (Mediterranean)

- *16. *Hormogaster redii* Rosa, 1887 [single record of transport to North America (Gates, 1954; 1972: 61)].

Family LUTODRILIDAE McMahan, 1976

- 17. *Lutodrilus multivesiculatus* McMahan, 1976.

Family LUMBRICIDAE (Holarctic).

- *18. *Allolobophora chlorotica* (Savigny, 1826)

- *19. *Allolobophoridella eiseni* (Levisen, 1884)
- *20. *Aporrectodea caliginosa* (Savigny, 1826) [includes *Aporrectodea turgida* (Eisen, 1873)].
- *21. *Aporrectodea icterica* (Savigny, 1826) [sometimes misspelt "icteria"].
- *22. *Aporrectodea limicola* (Michaelsen, 1890)
- *23. *Aporrectodea longa* (Ude, 1885)
- *24. *Aporrectodea rosea rosea* (Savigny, 1826) [many synonyms including *Ap. bowcrowensis* Reynolds & Clapperton, 1996: 77 (this latter synonymy by Csuzdi & Zicsi (2003: 93)].
- *25. *Aporrectodea trapezoides* (Duges, 1828)
- *26. *Aporrectodea tuberculata* (Eisen, 1874)
- 27. *Bimastos gieseleri gieseleri* (Ude, 1895) Savannah, Georgia (sometimes put in synonymy of *B. tumidus*).
- 28. *Bimastos gieseleri hempeli* (Smith, 1915) (sometimes put in synonymy of *B. tumidus*).
- 29. *Bimastos heimbürgeri* Smith, 1928
- 30. *Bimastos lawrenceae* Fender, 1994 in Fender, McKey-Fender & Marshall, 1994 [an endemic lumbricid, known only from Vancouver Island, BC, Canada].
- 31. *Bimastos longicinctus* (Smith & Gittins, 1915) [cf. *B. parvus*].
- 32. *Bimastos palustris* H.F. Moore, 1895 (*Bimastos* sp. H.F. Moore, 1893) [sometimes attributed to *Allolobophora*, also mislisted as "J.P. Moore" by James (1995: 43)].
- *-33. *Bimastos parvus* Eisen, 1974 [syns. *beddardi* Michaelsen, 1894 (non Ribaucourt, 1896 = *Ap. caliginosa/trapezoides* complex); *parva udei* Ribaucourt, 1896 (non Sapkarev, 1972 = *Serbiona joncesapkarovi* Blakemore, 2004: 78); *consticta geminata* Friend, 1897; *?longicinctus* Smith & Gittins, 1915: 548 - some authors maintain this parthenogenetic taxon on the grounds that it has >60-64 partitions in the calciferous glands vs. 40-45 in *B. parvus* (S. James, pers. com. 7th Dec, 2004)]. [Note: recently transferred from *Allolobophoridella parva* to *Bimastos parvus* by Csuzdi & Zicsi (2003: 69, 71), although sometimes still reported as *Allolobophora parva* or even *Eisenia parva*, some authors revoke the probable *beddardi* and possible *longicinctus* synonyms. Regarded as a North American endemic genus, but this taxon commonly transported thus subject to relocation in USA and noted as an introduction, at least to Yukon Territory, Canada].
- 34. *Bimastos tumidus* (Eisen, 1874) [syn. *Bimastos ducis* Stephenson, 1933, although James (1995: 34) apparently attempts to maintain it]. Also reported from Mexico (Fragoso pers. comm. Sept. 2006).
- 35. *Bimastos welchi* (Smith, 1917)
- 36. *Bimastos zeteki* (Smith & Gittins, 1915) (sometimes misspelt "zeteski" or "zetzki").
- *37. *Dendrobaena attemsi* Michaelsen, 1902

- *38. *Dendrobaena hortensis* (Michaelsen, 1890) [several synonyms].
- *39. *Dendrobaena octaedra octaedra* (Savigny, 1826) [several synonyms].
- *40. *Dendrobaena pygmaea* (Savigny, 1826) [syns. *Allolobophora minima* Rosa, 1884 non Muldal, 1953 (= *Murchieona minuscula*); *cognettii* Michaelsen, 1903 nom. nov. pro *ribaucourti* Cognetti, 1901 non Bretscher, 1901 (= *L. rubellus*); *pygmaea cognettii*: Bouché, 1972].
- *41. *Dendrobaena veneta veneta* (Rosa, 1886: 674) [syns *veneta veneta* (Rosa, 1886: 674); *caucasica* Kulagin, 1889; *bogdanowii* Kulagin, 1889; *veneta zebra* Michaelsen, 1902; *veneta succinta* Rosa, 1905; *venetus concolor* Michaelsen, 1909; *veneta picta* Michaelsen, 1910; *veneta tumida* Friend, 1927 (non Eisen, 1874); *austriaca* Michaelsen, 1936; *veneta crassa* Malevics, 1947 [non Michaelsen, 1900]; *veneta minuta* Malevics, 1947; *svetlovia* Grieb, 1948)].
- ?*42. *Dendrodrilus rubidus norvegicus* (Eisen, 1874) [possibly a morph rather than subspecies; a tentative report from Michigan and Indiana by Smith (1917: 177)].
- #43. *Dendrodrilus rubidus rubidus* (Savigny, 1826) [many synonyms].
- ?*44. *Dendrodrilus rubidus subrubicundus* (Eisen, 1874) [possibly a morph rather than subspecies].
- ?*45. *Dendrodrilus rubidus tenuis* (Savigny, 1826) [possibly a morph rather than subspecies].
- ?*46. *Eisenia andrei* Bouché, 1972 [syns. *E. fetida andrei* Bouché, 1972 replacement name for *E. fetida* var. *unicolor* André, 1963 as varietal names were invalid after 1961. Note: *E. fetida andrei*, placed in synonymy of *E. fetida* by Easton (1983) and Csuzdi & Zicsi (2003: 143), is sometimes given separate specific status, but this is almost arbitrary between authors - see discussion in Blakemore (2002)].
- *47. *Eisenia fetida* (Savigny, 1826) [many synonyms].
- ?*48. *Eiseniella tetraedra pupa* (Eisen, 1874) [syn. *hercynius* Michaelsen, 1890; *tetraedra quadripora* Cernosvitov, 1942. Recorded from Niagra; Germany; Portugal; USA (California and Illinois) and UK. Possibly meriting only varietal status as synonym of *Ei. t. tetraedra*, or maintained as a subspecies by some authors].
- *49. *Eiseniella tetraedra tetraedra* (Savigny, 1826) [many synonyms possibly including ?*Ei. tetraedra pupa* (Eisen, 1874) with syns. *hercynius* Michaelsen, 1890; *tetraedra quadripora* Cernosvitov, 1942 - according to Csuzdi & Zicsi (2003: 153) as the only differences of several *tetraedra* morphs are the locations of the male pores they could be either varieties rather than subspecies].
- 50. *Eisenoides carolinensis* Michaelsen, 1910 (syn. *pearsei* Stephenson, 1932).
- 51. *Eisenoides lonnbergi* (Michaelsen, 1894: 179) [originally *lönnerbergi*, sometimes spelt "loennerbergi" but, as the name is of Scandinavian rather than German origin (named for collector Dr Einar Lönnberg), under ICZN (1999: Art: 32.5.2.1) it is corrected with only the diatritic removed rather than changed to "oe"].

- *52. *Lumbricus castaneus* (Savigny, 1826)
- *53. *Lumbricus festivus* (Savigny, 1826)
- *54. *Lumbricus friendi* Cognetti, 1904 [in Washington-Baltimore Metropolitan Area (Csuzdi & Szlávecz, 2003)].
- *55. *Lumbricus rubellus* Hoffmeister, 1843
- *56. *Lumbricus terrestris* Linnaeus, 1758
- *57. *Murchieona muldali* (Omodeo, 1956) [nom. nov. pro *Allolobophora minima* Muldal, 1952 non *A. minima* Rosa, 1884 (= *Dendrobaena pygmaea*); the fate of *Allolobophora muldali* var. *pickfordi* Bouché, 1972 illegitimate varietal name is unknown. Note: *Mu. muldali* was resurrected from synonymy in *Mu. minuscula* Rosa, 1906 by Zicsi & Csuzdi (1999: 990)].
- *58. *Octolasion cyaneum* (Savigny, 1826)
- *59. *Octolasion lacteum* (Örley, 1881) [described from Ohio by Olsen (1933)].
- *60. *Octolasion tyrtaeum* (Savigny, 1826)
- *61. *Satchellius mammalis* (Savigny, 1826)

Family OCNERODRILIDAE

- *62. *Eukerria saltensis* (Beddard, 1892)
- #63. *Ocnerodrilus occidentalis occidentalis* Eisen, 1878 [from Fresno; syns: *O. o.* var. *sinensis* Eisen, 1900; *O. tenellulus* Gates, 1945].
- ?#64. *Ocnerodrilus occidentalis arizonae* Eisen, 1900:116 [from Phoenix: possibly synonymous with *O. occidentalis*, or with one of its pathenogenetic morphs].
- ?#65. *Ocnerodrilus mexicanus hawaiiensis* Eisen, 1900:124 [from San Francisco, brought in from Honolulu, Hawaii: possibly synonymous with *O. occidentalis*, or with one of its pathenogenetic morphs].

Family ACANTHODRILIDAE

[Records of *Diplocardia* may overlap Mexican borders but no species classed as a Mexican native by Fragoso *et al.* (1995: 111, Appendix) are listed here; the genus is defined by two gizzards in 5 and 6, seldom 6 and 7 (e.g. Michaelsen, 1900: 324; Stephenson, 1930: 850; 1933); cf. James (1995: 33) who states that there is but a single gizzard in *Diplocardia* (but extending through two segments). The genus may belong in a revived and redefined (sub)family Diplocardi-inae/-idae].

Acanthodrilidae taxa, that seem to have been overlooked in all the above references (except for Michaelsen, 1900), are from the genus *Deltania* Eisen, 1893: 250 (non *Deltacanthus* Diaz-Ungria & Rodrigo 1958: 42. n.n. pro *Deltania* Diaz-Ungria & Rodrigo 1957: 12) that is listed as a homonym

[on nomenclator <http://uio.mbl.edu/NZ/detail.php?uid=55691&d=1>], but was placed in synonymy of *Microscolex* Rosa, 1887 in Michaelsen (1900), viz.

Deltania benhami Eisen, 1893: 253 (= *Microscolex benhami*).

Deltania dubia Eisen, 1894: 22 (= a junior homonym and junior synonym of *M. dubius* (Fletcher, 1887) according to Michaelsen (1900) - from California).

Deltania elegans Eisen, 1893: 251 [= *Microscolex elegans* - from California; sometimes misdated "Eisen, 1892" e.g. Reynolds & Cook (1976)].

Deltania troyeri Eisen, 1893: 251 (?= *Microscolex troyeri troyeri*).

Deltania troyeri crassa Eisen, 1896: 169 (= *Microscolex troyeri*)

Deltania troyeri lagunae Eisen, 1896: 170 (= *Microscolex troyeri*)

-66. *Diplocardia alabamiana* Gates, 1977

-67. *Diplocardia alba alba* Gates, 1943: 108 [from Fort Myers, Florida; other subspecies *D. alba mexicana* Gates, 1955: 229 and *D. alba gravida* Gates, 1977, overlooked or synonyms from Mexico (omitted from Fragoso *et al.*, 1995: 111, Appendix). Cf. *Eodrillus albidus* Gates, 1973: 22 (= ?*Microscolex*) - nom. nov. pro *Eodrillus albus* Gates, 1970: 267 (non *Diplocardia alba* Gates, 1943: 108) from a cave in Mexico cited as "*Diplostrema albidus* .. Gates, 1970b, 1973" by Fragoso *et al.*, (1995: 111); non *Trigaster albida* Gates, 1973: 22 (= *Zapatadrillus albidus*) also from Mexico described in the same paper - this species was cited as "*albida*" in Reynolds & Cook (1976: 67), but miscited as "*T. alba* (see Gates, 1971; cf. *Zapatadrillus*)" in James (1991: 339), as "*Trigaster alba* Gates, 1973" in James (1991: 349), and as "*Z. albus*" or "*Zapatadrillus alba*" (sic) in James (1991: 347, 350, 352); finally corrected to *Zapatadrillus albidus* (Gates, 1973) in Fragoso *et al.*, (1995: 114); nec *Acanthodrillus albus* Beddard, 1895 (?= *Notiodrillus* or ?*Microscolex*) from Chile].

-68. *Diplocardia biprostatica* Gates, 1977

-69. *Diplocardia bitheca* Gates, 1977

-70. *Diplocardia bivesiculata* Murchie, 1961

-71. *Diplocardia californiana* James, 1994

-72. *Diplocardia caroliniana* Eisen, 1899 [originally *Diplocardia singularis caroliniana*].

-73. *Diplocardia communis communis* Garman, 1888 (sometimes misspelt "Garmann" or "Garmon").

?-74. *Diplocardia communis wolcotti* Macnab & McKey-Fender, 1955 [omitted by James (1990; 1995); type material in the authors' collection].

-75. *Diplocardia conoyeri* Murchie, 1961

?-76. *Diplocardia egglestoni* Murchie, 1958 [omitted by James (1990; 1995), description cited

online here: https://kb.osu.edu/dspace/bitstream/1811/4559/1/V58N05_270.pdf].

- #77. *Diplocardia eiseni* (Michaelsen, 1894) (originally *Geodrilus eiseni* from Florida, also known from Mexico).
- 78. *Diplocardia farmvillensis* Gates, 1977
- 78. *Diplocardia floridana* F. Smith, 1924
- 80. *Diplocardia fusca* Gates, 1943 [from Fort Worth, Texas].
- 81. *Diplocardia fuscula* Gates, 1968
- 82. *Diplocardia gatesi* Murchie, 1965
- 83. *Diplocardia glabra* Gates, 1967
- 84. *Diplocardia gracilis* Gates, 1943 [from Tennessee].
- 85. *Diplocardia hulberti* James, 1988
- #86. *Diplocardia invecta* Gates, 1955 (from Texas and Mexico)
- 87. *Diplocardia kansensis* James, 1990
- #88. *Diplocardia keyesi* (Eisen, 1896) [originally from Ensenada, Baja California, Mexico; listed in Table of "The currently recognized names of North American earthworms from the area east of the Rocky Mountains" by James (1995: 34), i.e. from USA, but possibly a mistake and it is confined to Mexico; two or three other similar Californian taxa mentioned by Wood & James (1993) here: <http://www.mycorrhiza.org/worms.pdf> were described by James (1994); originally *Aleodrilus keyesi* the type of Eisen's (1896) genus].
- #89. *Diplocardia koebeli* Eisen, 1900: 197 [correction (by Michaelsen, 1900: 325) of "koebeli" from Mexico and listed by James (1990: 383) under Chapter heading "Oligochaeta: Megascolecidae and other earthworm from southern and midwestern North America" i.e. from USA, but possibly this is a mistake and it is confined to Mexico; this was type species of Eisen's (1900) subgenus *Naillenia*, more recently transferred to *Protozapotecia*].
- 90. *Diplocardia komareki* Gates, 1977
- 91. *Diplocardia longa* J. Moore, 1905
- 92. *Diplocardia longiseta* Murchie, 1963
- 93. *Diplocardia macdowellii* Murchie, 1967 [compared to *D. michaelseni*].
- 94. *Diplocardia meansi* Gates, 1977
- 95. *Diplocardia michaelseni* Eisen, 1899 [possibly has non-tubular prostates; redescribed by F. Smith (1923) and by Stephenson (1933: 930-934) as "not obviously tubular; each appears to be essentially a rather loose elongated mass irregularly bent on itself, but even when unravelled as far as possible and its bends displayed it does not form a definite cylindrical "tube""].

- 96. *Diplocardia minima* Gates, 1977
- 97. *Diplocardia mississippiensis* F. Smith, 1924
- 98. *Diplocardia montana* James, 1994
- 99. *Diplocardia nova* Gates, 1977
- 100. *Diplocardia ornata* Gates, 1943 [from Tennessee].
- 101. *Diplocardia patuxentis* Csuzdi & Szlavecz, 2002
- 102. *Diplocardia pettibonae* Gates, 1977
- 103. *Diplocardia riparia riparia* F. Smith, 1895
- ?-104. *Diplocardia riparia prosenderis* Macnab & McKey-Fender, 1955 [possibly a junior synonym of the nominal sub-species, or (as by James, 1990: 384) listed at species level; unrepresented by type material which is listed in the authors' collection].
- 105. *Diplocardia rugosa* James, 1988
- 106. *Diplocardia sandersi* Gates, 1955
- ?-107. *Diplocardia singularis fluviatilis* F. Smith, 1915 [omitted by James (1990)].
- 108. *Diplocardia singularis singularis* (Ude, 1893) [originally *Geodrilus singularis* from Danville, Kentucky, described as *D. communis singularis* (Ude) by Michaelsen (1900: 326)].
- 109. *Diplocardia smithii* MacNab & McKey-Fender, 1955
- 110. *Diplocardia sylvicola* Gates, 1977
- 111. *Diplocardia texensis* F. Smith, 1924 [originally *Diplocardia keyesi* var. *texensis*].
- 112. *Diplocardia udei* Eisen, 1899 [possibly has non-tubular prostates and thus meriting transfer to another genus; not "*D. udei* Gates, 1955" as stated by James (1995: 34)].
- 113. *Diplocardia vaili* Gates, 1977
- 114. *Diplocardia varivesicula* Murchie, 1966 (often misspelt "varivesiculata").
- ?-115. *Diplocardia verrucosa recta* Murchie, 1962 [omitted by James (1990)].
- 116. *Diplocardia verrucosa verrucosa* Ude, 1895 [type of Ude's (1900) subgenus *Omahania*, from Nebraska, Omaha and also recorded from New Mexico state, USA]
- 117. *Diplocardia woodi* James, 1994
- 118. *Microscolex benhami* (Eisen, 1893) - from Alameda County, California.
- *119. *Microscolex dubius* (Fletcher, 1887) (synonyms include *Deltania dubia* Eisen, 1894, and *M. carolinae* Eisen, 1900).
- 120. *Microscolex elegans* (Eisen, 1893) - from Golden Gate Park, San Francisco, California.
- *121. *Microscolex phosphoreus* (Dugès, 1837) [synonyms probably include *Microscolex hempeli* F. Smith, 1896 and its synonyms: *M. parvus* + *M. parvus carolinianus* Eisen, 1900].
- 122. *Microscolex troyeri* (Eisen, 1893) (originally *Deltania troyeri* from Golden Gate Park, San Francisco; synonyms, *Deltania troyeri crassa* + *Deltania troyeri lagunae* Eisen, 1896); possibly a synonym of *M. phosphoreus*.

Family OCTOCHAETIDAE (Benhamiinae)

- *123. *Dichogaster (Diplotheodrilus) affinis* (Michaelsen, 1890) [From Florida and Galapagos by Csuzdi (1997: 37; 2000: 59)].
- *124. *Dichogaster (Diplotheodrilus) bolau* (Michaelsen, 1891)
- *125. *Dichogaster (Diplotheodrilus) modiglianii* (Rosa, 1896) [From Florida by Csuzdi (1997)].
- *126. *Dichogaster (Diplotheodrilus) saliens* (Beddard 1893)

Family MEGASCOLECIDAE

(natives allocated in tribe Argilophilini after Fender & McKey-Fender, 1990: 370)

- 127. *Arctiostrotus adunatus* McKey *et al.* in Fender, McKey-Fender & Marshall, 1994
- 128. *Arctiostrotus altmani* (Gates, 1942) [sometimes cited as Macnab & McKey Fender, 1948 due to mistake in Reynolds & Cook (1976); replacement name for *Plutellus decathecus* Altman, 1936 non Michaelsen, 1910 (= *Vesiculodrilus decathecus*) from Tasmania].
- 129. *Arctiostrotus fontinalis* McKey *et al.* in Fender, McKey-Fender & Marshall, 1994
- 130. *Arctiostrotus johnsoni* McKey *et al.* in Fender, McKey-Fender & Marshall, 1994
- 131. *Arctiostrotus perrieri* (Benham, 1892)
- 132. *Arctiostrotus pluvialis* McKey *et al.* in Fender, McKey-Fender & Marshall, 1994
- 133. *Arctiostrotus vancouverensis* McKey *et al.* in Fender, McKey-Fender & Marshall, 1994
- 134. *Argilophilus collinus* Eisen, 1900 (originally *Argilophilus marmoratus collinus*).
- 135. *Argilophilus garloughi* (F. Smith, 1937)
- 136. *Argilophilus hammondi* (McKey-Fender, 1970: 235)
- 137. *Argilophilus hyalinus* Eisen, 1900
- 138. *Argilophilus margaritae* James, 1994
- 139. *Argilophilus marmoratus marmoratus* Eisen, 1893
- ?-140. *Argilophilus marmoratus ornatus* Eisen, 1893; Sebastopol, California
- 141. *Argilophilus panulirus* (MacNab & McKey-Fender, 1959)
- 142. *Argilophilus papillifer* Eisen, 1893 (originally *Argilophilus marmoratus papillifer*); Berkeley, California.
- 143. *Argilophilus sierrae* (Michaelsen, 1921)
- 144. *Argilophilus woodi* James, 1994
- 145. *Chetcodrilus exutus* (MacNab & McKey-Fender, 1958: 107) [originally *Plutellus fenderi exutus*].
- 146. *Chetcodrilus fenderi* (MacNab in MacNab & McKey-Fender, 1958: 103)
- 147. *Chetcodrilus umbellulariae* (MacNab & McKey-Fender, 1958)
- 148. *Drilochaera chenowithensis* (McKey-Fender, 1970: 240)

- 149. *Driloleirus americanus* (F. Smith, 1897)
- 150. *Driloleirus cascadenis* (F. Smith, 1937)
- 151. *Driloleirus eiseni* (F. Smith, 1937)
- 152. *Driloleirus macelfreshi* (F. Smith, 1937)
- 153. *Driloleirus michaelseni* (F. Smith, 1937)
- 154. *Driloleirus wellsi* (Altman, 1936)
- 155. *Kincaidodrilus kincaidii* (Altman, 1936)
- 156. *Macnabodrilus hopsonae* (McKey-Fender, 1970: 225) [Reynolds & Cook (1992: 9) miscite this taxon as the type of the genus (cf. *M. macnabi*), they also misdate it as "1957"].
- 157. *Macnabodrilus macnabi* (McKey-Fender, 1957) [type of genus].
- 158. *Nephrellaxis blacki* (MacNab & McKey-Fender, 1953)
- 159. *Nephrellaxis davisii* (MacNab & McKey-Fender, 1953)
- 160. *Toutellus adecus* (MacNab & McKey-Fender, 1959)
- 161. *Toutellus hyoiedes* (MacNab & McKey-Fender, 1959) (misspelt "hyoides" in Reynolds & Cook, 1992: 16).
- 162. *Toutellus oregonensis* (F. Smith, 1937)
- 163. *Toutellus toutellus* (Altman, 1936)
- *164. *Amyntas agrestis* (Goto & Hatai, 1899) [introduced into North America eg. reported from New York, Connecticut and Louisiana by Gates (1958: 1) and by Gates (1982) who thought it imported from Japan; also from Georgia (Callahan *et al.*, 2004); and North and South Carolina and Tennessee (Wetzel, 2005). (Syn. *Pheretima striata* Ishizuka, 1999, for fuller details - see Blakemore, 2003, 2005)].
- *165. *Amyntas corticis* (Kinberg, 1867) [many synonyms including *Megascolex diffringens* Baird, 1869].
- *166. *Amyntas gracilis* (Kinberg, 1867) [many synonyms including *Perichaeta hawayana* Rosa, 1891].
- *167. *Amyntas hupeiensis* (Michaelsen, 1895) (first recorded in Washington, D.C. in 1910, since spread to other States, eg. New York, Maine, Connecticut, Florida, Philadelphia, Illinois, North and South Carolina, Tennessee). Gates (1958: 17-20) thought American worms in greenhouses, (?turf farms), and golf courses may have been introduced from Japan.
- *168. *Amyntas loveridgei* (Gates, 1968)
- *169. *Amyntas minimus* (Horst, 1893)
- *170. *Amyntas morrissi* (Beddard, 1892)
- *171. *Amyntas rodericensis* (Grube, 1879)
- *172. *Amyntas tokioensis* (Beddard, 1892) **new US record from synonyms.** [Syns. reported from Schenectady, New York City, and New Jersey as *Pheretima levis* (Goto & Hatai, 1899)

by Gates, 1958: 21-24 (?syns. *irregularis*, *ambigua*, *vittata*, *schizopora*) by Gates (1958: 21-22) or as *Metaphire levis* (Goto & Hatai, 1899) by Easton (1981: 53); although incorrectly cited as “*Metaphire levis* (Horst, 1893)” from North Carolina and Tennessee, USA by Wetzel (2005), and as *Amynthas vittatus* (Goto & Hatai, 1898) from Ashford, northeastern Connecticut by Schneider & McDevit (2002); - for comb. nov. and full synonymy see Blakemore (2003, 2005)].

*173. *Metaphire californica* (Kinberg, 1867)

*174. *Metaphire hilgendorfi* (Michaelsen, 1892) [reported from Louisiana and Middleburgh, Virginia by Gates (1958: 11-12) and from South Carolina and Tennessee by Wetzel (2005); for numerous and full synonymy - see Blakemore (2003, 2005)].

*175. *Metaphire houletti* (Perrier, 1872)

*176. *Metaphire posthuma* (Vaillant, 1868)

*177. *Perionyx excavatus* Perrier, 1872

*178. *Pithemera bicincta* (Perrier, 1875)

*179. *Polypheretima elongata* (Perrier, 1872)

*180. *Pontodrilus litoralis* (Grube, 1855)

Family GLOSSOSCOLECIDAE

*181. *Pontoscolex corethrurus* (Müller, 1856)

Family EUDRILIDAE

*182. *Eudrilus eugeniae* (Kinberg, 1867)

Species dubius

Hypogaeanum hirtum Savigny, 1820: 104 terrestrial from Pennsylvania (Philadelphia), listed as *species dubius* in Michaelsen (1900: 519); types missing.

Lumbricus apii Kinberg, 1867:100 described from Sausalita Bay, San Francisco but abandoned after Michaelsen's (1900: 520) *species dubius* listing; types are in Leiden: 1939.

References

- Blakemore, R.J. (2000). *Tasmanian Earthworms - with Review of World Families*. VermEcology, Kippax, Canberra. Pp. 800 including 222 figures (CD format). Published December, 2000.
- Blakemore, R.J., (2002). *Cosmopolitan Earthworms – an Eco-Taxonomic Guide to the Peregrine Species of the World*. VermEcology, Kippax, Australia. Pp. 426 + 80 figs (CD format).

- Blakemore, R.J., (2003). Japanese Earthworms (Annelida: Oligochaeta): a Review and Checklist of Species. *Organisms, Diversity and Evolution*. 3(3): 241-244. [Published Sept., 2003].
<http://www.urbanfischer.de/journals/ode/> + Electronic Supplement 2003-11.
<http://www.senckenberg.de/odes/03-11.htm> [Published after October, 2003].
- Blakemore, R.J. (2005). *A Series of Searchable Texts on Earthworm Biodiversity, Ecology and Systematics from Various Regions of the World*. Eds.: N. Kaneko & M.T. Ito. COE Soil Ecology Research Group, Yokohama National University, Japan. CD-ROM.
<http://bio-eco.eis.ynu.ac.jp/eng/database/earthworm/> July, 2005].
- Blakemore, R.J., (2006). *Cosmopolitan Earthworms – an Eco-Taxonomic Guide to the Peregrine Species of the World*. Volume 2. VermEcology. Pp. 600 + 100 figs (CD format).
- Callahan, M.A., Jr., Hendrix, P.F. & Phillips, R.J. (2003). Population Dynamics of an exotic earthworm (*Amyntas agrestis*) in undisturbed soils of the southern Appalachian Mountains, USA. *Pedobiologia* 47:466-470.
- Csuzdi, Cs. & Szilávecz, K. (2003). *Lumbricus friendi* Cognetti, 1904 a new exotic earthworm in North America. *Northeastern Naturalist* **10(1)**: 77-82.
- Edwards, C.A. & Lofty, J.R. (1977). "Biology of Earthworms." (2nd edn.). Chapman and Hall, London. Pp. 333.
- Fender, W. M., (1995). Native earthworms of the Pacific Northwest: an ecological overview. Pages 53-66 in P. F. Hendrix (ed.). "Ecology and biogeography of earthworms in North America". CRC Publishing, Boca Raton, Florida.
- Fender, W. M., & McKey-Fender, D. (1990). Oligochaeta: Megascolecidae and other earthworms from western North America. Pages 357-378, in D. L. Dindal (ed.). "Soil biology Guide". John Wiley & Sons, Inc., New York.
- Gates, G.E. (1942). Check List and Bibliography of North American Earthworms. *American Midland Naturalist*. **27(1)**: 86-108.
[http://links.jstor.org/sici?sici=0003-0031\(194201\)27%3A1%3C86%3ACLABON%3E2.0.CO%3B2-K](http://links.jstor.org/sici?sici=0003-0031(194201)27%3A1%3C86%3ACLABON%3E2.0.CO%3B2-K)].
- Gates, G.E. (1958). On some species of the oriental earthworm genus *Pheretima* Kinberg, 1867, with a key to species reported from the Americas. *American Mus. Novitates*. 1888: 1-33.
<http://digitallibrary.amnh.org/dspace/handle/2246/4574>].
- Gates, G.E. (1972). Burmese Earthworms, an introduction to the systematics and biology of Megadrile oligochaetes with special reference to South-East Asia. *Transactions of the American Philosophical Society*. **62(7)**: 1-326. [Online via subscription here
<http://links.jstor.org/journals/00659746.html>].

- Gates, G.E. (1972a). Contributions to North American earthworms (Annelida: Oligochaeta). No. 3: Towards a revision of the earthworm family Lumbricidae IV. The *trapezoides* species group. *Bulletin of the Tall Timbers Research Station*. **12**: 1-146.
- Gates, G.E. (1982). Farewell to North American megadriles. *Megadrilogica* 4(1-2): 12-77.
- James, S.W. (1990). Oligochaeta: Megascolecidae and other earthworms from southern and midwestern North America. Pages 379-386, in D.L. Dindal (ed.). "Soil Biology Guide". John Wiley and Sons, New York.
- James, S.W. (1995). Systematics, biogeography and ecology of earthworms from eastern, central, southern and southwestern USA. in P. Hendrix (ed.). "Earthworm Ecology and Biogeography in North America". CRC Press, Inc, Boca Raton, Florida. Pp. 29-51.
- Fragoso, C., James, S.W., & Borges, S. (1995). Native earthworms of the north. Neotropical region: current status and controversies. in P. Hendrix (ed.) Earthworm Ecology and Biogeography in North America. CRC Press, Inc, Boca Raton, Florida. Pp. 67-115.
- Hendrix, P.F & Bohlen, P.J. (2002) Exotic earthworm invasions in North America: Ecological and policy implications. *Bioscience*. 52(9): 801-811. [May, 2005, URL: www.eeb.uconn.edu/Courses/Eeb302/Bioscience2002BohlenEarthwormsPolicy.pdf].
- Michaelsen, W. (1900). Das Tierreich 10: Vermes. Friedman & Sohn. Germany. Pp. 715.
- Murchie, W.R. (1967). Description of *Diplocardia Macdowelli* a New Megascolecid Earthworm from Mississippi (Oligochaeta). *The Ohio Journal of Science*. 67(1): 50-53. [https://kb.osu.edu/dspace/bitstream/1811/5275/1/V67N01_050.pdf].
- Olsen, H. W. (1933). Two New Species of Earthworms for Ohio. *The Ohio Journal of Science*. **33(3)**: 192-193. [Online: https://kb.osu.edu/dspace/bitstream/1811/2630/1/V33N03_192.pdf].
- Reynolds, J.W. (1977). The earthworms (Lumbricidae and Sparganophilidae) of Ontario. *Life sciences Miscellaneous Publications*, Royal Ontario Museum, Canada.
- Reynolds, J.W. & Wetzel, M.J. (2004). Terrestrial Oligochaeta (Annelida: Clitellata) in North America north of Mexico. *Megadrilogica* 9(11): 71-98. [March, 2004].
- Schneider, C.W. & McDevit, D.C. (2002). Are earthworms a possible mechanism for airborne dispersal of the alga *Vaucheria*? *Northeastern Naturalist* **9(2)**: 225-234.
- Schwert, D.P. (1990). Oligochaeta: Lumbricidae. Pages 341-356, in D. L. Dindal, (ed.). "Soil Biology Guide". John Wiley & Sons, Inc., New York.
- Sims, R.W. and Gerard, B.M. (1985). Earthworms. Keys and notes to the identification and study of the Species. *Synopsis of the British Fauna (New series)*. Brill, Leiden. No. **31**. Pp. 171.
- Sims, R. W. & Gerard, B.M. (1999). *Earthworms: Notes for the identification of British species*. 4th Edition. The Linnean Society of London and The Estuarine and Coastal Sciences

- Association by Field Studies Council, Montford Bridge, Shrewsbury, UK. Pp. 169.
- Smith, F. (1923). A Revision of the Description of *Diplocardia michaelsoni* Eisen. *Transactions of the American Microscopical Society*, 42(4): 175-179.
- Stephenson, J. (1930). *The Oligochaeta*. Oxford Press.
- Wetzel, M. J. (2003). The Terrestrial Oligochaeta of Illinois - annotated checklist of species. WWW URL: [<http://www.inhs.uiuc.edu/cbd/collections/annelid/ilspecies.html>] 12 March 2003.
- Wetzel, M. J. (2004). The Terrestrial Oligochaeta of Illinois - Annotated Checklist of Species. WWW URL: [<http://www.inhs.uiuc.edu/~mjwetzel/TerrWoi.mjw.list.html>]. 23 April 2004.
- Wetzel, M.J. (2005). Checklist of the Aquatic and Terrestrial Oligochaeta occurring in North Carolina, South Carolina, and / or Tennessee, with notations for species now known to occur in the Great Smoky Mountains National Park. WWW URL: [<http://www.inhs.uiuc.edu/~mjwetzel/AOGSMNP.PkChklist.html>] May, 2005; March, 2006.
- Wood, H.B. & S.W. James (1993). Native and introduced earthworms from selected chaparral, woodland, and riparian zones in southern California. *Gen. Tech. Rep. PSW-GTR-142*. Albany, CA: Pacific Southwest Research Station, Forest Service, USDA; Pp. 20.

[End of North American Checklist].